



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION BY A. M. HARDING, University of Arkansas.

$$f'(x) = -e^{-cx}(c + c \cos x + \sin x).$$

$$f''(x) = e^{-cx}(c^2 + c^2 \cos x + 2c \sin x - \cos x).$$

Now $f'(x) = 0$ only when $c + c \cos x + \sin x = 0$,
that is, when

$$c(1 + \cos x) + \sin x = 0,$$

or

$$2c \cos^2 \frac{x}{2} + 2 \sin \frac{x}{2} \cos \frac{x}{2} = 0.$$

Hence,

$$\cos \frac{x}{2} = 0, \quad \text{and} \quad x = \pi, 3\pi, 5\pi, \dots (2n-1)\pi, \dots$$

or

$$c \cos \frac{x}{2} + \sin \frac{x}{2} = 0, \quad \text{and} \quad x = 2 \arctan (-c).$$

When $x = \pi, 3\pi, 5\pi, \dots, (2n-1)\pi, \dots$, $f''(x) = -c(2n-1)\pi$, which is always positive.

Hence, the minimum value of $f(x)$ is obtained by giving x any of these values. When $x = 2 \arctan (-c)$, that is, $c \cos x/2 + \sin x/2 = 0$,

$$\begin{aligned} f''(x) &= e^{-cx}[c^2(1 + \cos x) + c \sin x - \cos x] \\ &= e^{-cx} \left[2c \cos \frac{x}{2} \left(c \cos \frac{x}{2} + \sin \frac{x}{2} \right) + 2c \sin \frac{x}{2} \cos \frac{x}{2} - \cos x \right] \\ &= e^{-cx} \left[2c \sin \frac{x}{2} \cos \frac{x}{2} - \cos x \right] = e^{-cx} \left[2 \sin \frac{x}{2} \left(-\sin \frac{x}{2} \right) - \cos x \right] \\ &= -e^{-cx} \left[2 \sin^2 \frac{x}{2} + \cos x \right] = -e^{-cx}[1 - \cos x + \cos x] = -e^{-cx}. \end{aligned}$$

Now $-e^{-cx}$ is negative for the above value of x . Hence the maximum value of $f(x)$ is obtained by giving x the value $2 \arctan (-c)$.

Also solved by W. C. EELLS, PAUL CAPRON, H. C. FEEMSTER, G. W. HARTWELL and the PROPOSER.

MECHANICS.


286. Proposed by C. N. SCHMALL, New York City.

A slightly elastic string is just long enough to reach between two hooks on the same horizontal line. A ring of weight w is placed at its middle point. Show that the ring will sink through a distance $h = a\sqrt{3e\omega/2}$, where e is the elasticity of the string and $2a$ the distance between the two hooks.

SOLUTION BY B. F. FINKEL, Drury College.

Since w is at the middle point of the string, the tension T in the two halves of the string is the same when the string is in equilibrium. Let θ be the angle which the string makes with the horizontal line.

Then $AC = a \sec \theta$. Hence the strain in AC is



$$\frac{a \sec \theta - a}{a} = \sec \theta - 1.$$

By Hooke's Law, we have $T/(\sec \theta - 1) = e$. (1)

For equilibrium, we have $2T \sin \theta = w$. (2)

Hence,

$$2e \sin \theta (\sec \theta - 1) = w,$$

or

$$\tan \theta - \sin \theta = \frac{w}{2e}. \quad (3)$$

Since the string is only slightly elastic, θ must be small.

When θ is small we may take $\tan \theta = \theta + (\theta^3/3)$ and $\sin \theta = \theta - (\theta^3/6)$. Hence, substituting these values in (3) and solving for θ , we have $\theta = \sqrt[3]{w/e}$.

But $\tan \theta = h/a$, or $\theta + (\theta^3/3) = h/a$. Substituting for θ and solving for h , we have

$$h = a \left\{ \sqrt[3]{\frac{w}{e}} + \frac{w}{3e} \right\}.$$

The proposer's result is incorrect, as the test of physical dimensions shows.

290. Proposed by B. F. FINKEL, Drury College.

A fox, pursued by a hound, is running with uniform velocity over a frail arch in the form of a cycloid: the hound stops at a weak point of the arch, then tumbles through, and reaches the level ground with a velocity equal to that of the fox. Prove that the fox exerted no normal pressure on the arch at the point where the hound fell through. Walton's *Problems in Theoretical Mechanics*, p. 662.

SOLUTION BY THE PROPOSER.

Let $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$ be the parametric equations of the cycloidal arch and v_0 , the speed of the fox. Then at the point where the hound tumbled through, we have the relation $v_0^2 = 2gy_0$, where y_0 is the ordinate of the arch at that point.

Now the centripetal force of the fox is $m(v_0^2/R)$, where m is his mass and R the radius of curvature of the arch. The force exerted by gravity in opposition to this force is $mg \cos \phi$, where ϕ is the direction of the arch at any point.

Hence, the normal pressure on the arch at any point, assuming $m = 1$, is

$$P = \frac{v_0^2}{R} - g \cos \phi.$$

From the equations of the arch, we have

$$\frac{dy}{dx} = \tan \phi = \cot \frac{\theta}{2} = \tan \left(\frac{\pi}{2} - \frac{\theta}{2} \right).$$